Aquablanc

Aquablanc O₂ Gentle

Key Features/Characteristics

Gentler approach to pool care

Maintain pH levels within the ideal range of 7.0 - 7.4.

Application Instructions:

Before converting a pool to Aquablanc O₂ Gentle, shock dose with Blue Horizons Granular Shock, Rapid Shock or Shock 'n' Floc to a minimum of 5 mg/l but ideally 10mg/l free chlorine.

- 1. **Initial Dose:** Add two double sachets (2 x powder sachets & 2 liquid sachets) per 4,400 gallons (20m³) of pool water.
- 2. **Weekly Dose:** Add one double sachet (1 x powder sachet & 1 liquid sachet) per 4,400 gallons (20m³) of pool water.
- 3. Carefully apply the powder sachet first aiming for even distribution close to the pool water surface. Apply whilst the filter pump(s) are turned 'on' to aid distribution. **Warning:** When dosing this sachet, check wind direction first to ensure granules are not blown back towards you.
- 4. Pour the liquid sachet into the pool, whilst the pump(s) are turned 'on' preferably near the water inlets to aid distribution.
- 5. Test the oxygen level one hour after applying Aquablanc O₂ Gentle. The oxygen level should be between 5 8mg/l. If the oxygen figure is below this level, the weekly dose rate of O₂ Gentle may need to be increased. Alternatively it may be time to shock dose the swimming pool with Aquablanc Active Oxygen Liquid or a Blue Horizons chlorine shock product.

Usage Application						
m ¹ /gallons Initial 1st day	20 m ³ (4,400) 2x (3 B)	30 m³ (6,600) 3x	40 m³ (8,800) 4x	50 m ³ (11,000) 5x	60 m³ (13,200) 6x (13,200)	70 m ³ (15,400) 7x (10 B)
regularly every	7th day 1x	4th day	7th day 2x	5th day 2x	7th day 3x	6th day 3X

